

Wild dog risks to threatened wildlife

Wild dogs prey on a variety of animals including mammals, birds and reptiles of all sizes from insects to water buffalo. However, they prefer to eat small and medium-sized mammals when available, including native mice, dunnarts, bandicoots and wallabies¹.

Wild dogs can present significant risks to populations of many of these species², and the continual influx of domestic dogs into the wild means there is a constant feral population that puts pressure on vulnerable wildlife³.

Wild dogs have been implicated in the decline of several species, both historically and in the recent past. For example, dingoes may have been responsible for the mainland extinction of the thylacine (Tasmanian tiger)¹, as well as the decline of brushtail possums⁴ and other marsupials⁵ in the arid zone over the last 200 years.

Wild dogs have been implicated in the decline of several species, both historically and in the recent past.

Predation by wild dogs was recently listed as a Key Threatening Process for threatened species, populations and communities in New South Wales³ and is a known or potential threat to a variety of other native species elsewhere.

There are presently 14 national-level recovery plans that identify wild dogs as a known or potential threat to some native mammal, bird and reptile species (Table 1). These species are all listed under the national *Environment Protection and Biodiversity Conservation Act* 1999.



Image: Ben Allen

The recovery plans set out research and management actions that aim to stop the threatened species' decline and maximise their long-term survival in the wild. Several other plans are being prepared for the recovery of threatened species that would be potentially susceptible to predation by wild dogs should they re-establish in the places where these native species live.

Because the impact of wild dogs on populations of threatened species can be severe, the risk to all wildlife (threatened or otherwise) should be considered in regional wild dog management plans.







Table 1. Currently listed native species potentially or known to be threatened by wild dog predation⁶

Species type	Common name	Scientific name	Adult weight (kg)
mammal	marsupial moles	Notorycetes typhlops, N. caurinus	0.07
mammal	smoky mouse	Pseudomys fumeus	0.09
bird	black-breasted button-quail	Turnix melanogaster	0.1
mammal	golden bandicoot	Isoodon auratus	0.7
mammal	northern quoll	Dasyurus hallucatus	1.2
mammal	greater bilby	Macrotis lagotis	2.5
mammal	long-footed potoroo	Potorous longipes	2.5
bird	malleefowl	Leipoa ocellata	2.5
mammal	bridled nailtail wallaby	Onychogalea fraenata	8.0
mammal	Proserpine rock-wallaby	Petrogale persephone	8.8
mammal	koala	Phascolarctos cinereus	12
mammal	northern hairy-nosed wombat	Lasiorhinus krefftii	31
bird	southern cassowary	Casuarius casuarius johnsonii	60
reptile	marine turtles	various	-

More information

- 1. Corbett LK (2001). *The Dingo in Australia and Asia* (2nd edn). JB Books, South Australia.
- Coutts-Smith AJ, Mahon PS, Letnic M, and Downey PO (2007). <u>The Threat Posed by Pest Animals to Biodiversity in New South Wales</u>. Invasive Animals Cooperative Research Centre, Canberra.
- Major R (2009). <u>Predation and Hybridisation by Feral Dogs (Canis lupus familiaris)</u> <u>Key Threatening Process Listing</u>. New South Wales Department of Environment, Climate Change and Water, Sydney.
- 4. Kerle JA, Foulkes JN, Kimber RG, Papenfus D (1992). The decline of the brushtail possum, *Trichosurus vulpecula* (Kerr 1798), in arid Australia. *The Rangeland Journal* 14:107-127.
- Allen BL (2011). A comment on the distribution of historical and contemporary livestock grazing across Australia: Implications for using dingoes for biodiversity conservation. <u>Ecological Management and Restoration</u> 12:26-30.

6. Environment Protection and Biodiversity Conservation Act 1999. Australian Government. www.environment.gov.au/epbc/



Bridled nailtail wallaby. Image: Peter Fleming